


Sub  
Ab

- 5



10

- 15

3. The cyclic peptide of claim 2, wherein Z has a side chain comprising oxygen, nitrogen or carbon.
4. The cyclic peptide of claim 2, wherein the cyclic bond is a lactam or lactone bond.
5. The cyclic peptide of claim 1 or claim 2, wherein the cyclic peptide is capable of inhibiting the *agr* response.

6. The cyclic peptide of claim 2, wherein the cyclic peptide is capable of inhibiting the *agr* response.
7. The cyclic peptide of claim 1, wherein y is 4.
8. The cyclic peptide of claim 2, wherein y is 4.
- 5 9. The cyclic peptide of claim 7, wherein the peptide has an amino acid sequence that comprises G-V-N-A-X-S-S-L-F (Seq.ID No.:1), G-A-N-A-X-S-S-L-F (Seq.ID No.:2), G-V-A-A-X-S-S-L-F (Seq.ID No.:3), A-V-A-N-X-S-S-L-F (Seq.ID No.:4), G-V-N-A-X-A-S-L-F (Seq.ID No.:5), G-V-N-A-X-S-A-L-F (Seq.ID No.:6), G-V-N-A-X-S-S-A-F (Seq.ID No.:7) or X-S-S-L-F (Seq.ID No. 8).
- 10 10. The cyclic peptide of claim 8, wherein the peptide has an amino acid sequence that comprises G-V-N-A-X-S-S-L-F (Seq.ID No.:1), G-A-N-A-X-S-S-L-F (Seq.ID No.:2), G-V-A-A-X-S-S-L-F (Seq.ID No.:3), A-V-A-N-X-S-S-L-F (Seq.ID No.:4), G-V-N-A-X-A-S-L-F (Seq.ID No.:5), G-V-N-A-X-S-A-L-F (Seq.ID No.:6), G-V-N-A-X-S-S-A-F (Seq.ID No.:7) or X-S-S-L-F (Seq.ID No. 8).
- 15 11. A composition comprising the peptide of claim 1 and a carrier.
12. A composition comprising the peptide of claim 2 and a carrier.
13. A pharmaceutical composition comprising the peptide of claim 1 and a pharmaceutically acceptable carrier.
- 20 14. A pharmaceutical composition comprising the peptide of claim 2 and a pharmaceutically acceptable carrier.

FOIA b7 - D

- wherein X is selected from the group consisting of an amino acid, an amino acid analog, a peptidomimetic and a non-amide isostere. Z is selected from the group consisting of a synthetic amino acid and a biosynthetic amino acid, R is selected from the group consisting of oxygen, nitrogen and carbon, n is 0 to 10 and y is 1 to 10, said method comprising:

A. assembling the linear constituents of the peptide under preparation on

- a PEGA resin support to form a protected and bound peptide chain;
- B. treating the peptide chain of Step A to cause deprotection thereof;
- C. treating the deprotected peptide of Step B with buffer at a neutral pH for a period of time sufficient to cleave said peptide from said solid phase support and to form said cyclic peptide; and
- D. recovering said cyclic peptide.

20. A method for the preparation of a cyclic peptide in accordance with Claim 1, which comprises:

- A. assembling the linear peptide chain corresponding in composition to said cyclic peptide on to a solid phase resin support containing 3-mercapto-propionamide-polyethylene glycol-poly-(N<sub>1</sub>N<sub>4</sub>-dimethacrylamide)(HS-PEGA) to form a protected assembled peptide;
- B. treating the protected assembled peptide of Step A to deprotect said assembled peptide;
- C. treating the deprotected peptide of Step B with aqueous buffer at a pH of about 7.0 for a period of time sufficient to form said cyclic peptide and to cleave said peptide from said solid phase resin support; and
- D. recovering said cyclic peptide.

21. The method of Claim 19, wherein said solid phase resin support comprises BOC - AA- (linear assembled peptide)-PEGA.

22. The method of Claim 19, wherein the treatment of Step B is performed with HF for about 1 hour.

23. The method of Claim 19, wherein the treatment of Step C is performed with a buffer comprising Na<sub>2</sub>PO<sub>4</sub> and acetonitrile.

24. The method of Claim 22, wherein the treatment of Step C is performed for about 12 hours.

ADD  
A8